

 **Total Label USA**
Proof Approval

PRODUCT NUMBER
440404

COLORS
BOOK



Cyan Magenta Yellow Black

BASE



Black

COMPANY

Westbridge Agricultural Products

LABEL DESCRIPTION

Earthtrend Soil Book

LABEL PART NUMBER

LABEL SIZE

5.5000" x 5.5000"

STOCK

White Bopp

FINISH

Gloss Lam

DATE

9/8/2021

BARCODE

ADDITIONAL INFORMATION

NOTE: Per my signature, I have read the information contained on this proof approval form including, but not limited to, copy, barcode number, color and size. I understand that my signature confirms that the information is my responsibility and is correct and labels will be produced according to this signed document.

Color proofs are not true representations of color. Use for color breaks only.

APPROVED

CHANGES

(please mark below)

Signature/Date

SEE FOLLOWING PAGES FOR ARTWORK

EarthTrend[®]



SOIL

PLANT GROWTH REGULATOR

Active Ingredient*:

Cytokinin (as kinetin)	00.004%**
Other Ingredients:	99.996%
TOTAL	100.000%

* Contains Non-Plant Food Ingredients ** Based on Biological Activity

KEEP OUT OF REACH OF CHILDREN
CAUTION

See inside booklet for FIRST AID, PRECAUTIONARY STATEMENTS, and DIRECTIONS FOR USE

SHAKE WELL BEFORE USING



Manufactured by:
SAN Group Biotech USA, Inc.
1260 Avenida Chelsea
Vista, CA 92081 USA

EPA REG. NO. 51517-3
EPA EST. NO. 51517-CA-01

Net Contents: 1 U.S. Gallon

PRODUCT INFORMATION

Earthtrend Soil is a plant growth regulator for use on agronomic and horticultural crops as listed on the label. Applied as a supplement to a cultural program at specific stages of plant development, **Earthtrend Soil** improves seedling vigor, crop quality, uniformity and growth. **Earthtrend Soil** reduces transplant shock in new plantings.

Use **Earthtrend Foliar** to:

- **Improve germination and growth** of small seeded crops (for example alfalfa, direct seeded onion, carrots and cucumbers).
- **Promote improved root growth** and nodulation in legumes.
- **Promote early seedling vigor** and the ability of a plant (for example bell peppers and tomatoes) to set and carry a crop to maturity.
- **Improve lateral tuberous root growth** (sweet potatoes), improve initiation growth and maturity of tubers and ornamental bulbs providing a more uniform crop, and increase specific gravity (potatoes) and number of tubers.
- **Promote rooting and growth** of ornamental varieties from slips or cuttings and bare root transplants.

Earthtrend Soil must be used as part of complete fertility program. **Earthtrend Soil** can be used in a program with **Earthtrend Foliar**. Please refer to **Earthtrend Foliar** label for application rates and timings.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID	
IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information call 1-800-876-2767 Monday-Friday 8-5 PT. After hours, weekends or holidays call 1-800-222-1222, the National Poison Control Center.	

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if inhaled or absorbed through skin. Avoid breathing spray mist and contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Water proof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions are given for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product.
- Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Treated seed exposed on soil surface may be hazardous to birds and other wildlife. Cover or collect seeds spilled during loading. Do not use treated seed for food, feed or oil purposes. Treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of all excess treated seed and seed packaging by burial away from streams and bodies of water. Only protected handlers may be in the area during application.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the state or tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this labeling about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Waterproof Gloves
- Shoes plus socks

FOR USE ON

FIELD CROPS: Alfalfa (Hay and Seed Production), Canola, Corn (including Popcorn), Cotton, Hops, Peanuts, Rice, Sorghum (Milo), Soybeans, Sugarcane, Wheat and other Cereal Grains.

FRUITS and NUTS: Avocados, Bananas and Plantains, Berry Fruits (such as Blueberries, Blackberries, Currants, Huckleberries, Loganberries and Raspberries), Citrus Fruits (such as Grapefruit, Lemons, Limes and Oranges), Coffee, Cranberries, Exotic Fruits (such as Figs, Guava, Kiwifruit, Mangos, Papayas, Persimmons and Pineapples), Grapes, Pome Fruits (such as Apples, Crabapple, Pears and Quinces), Stone Fruits (such as Apricots, Cherries, Peaches, Plums and Nectarines), Strawberries, Tree Nuts (such as Almonds, Cashews, Hazelnuts, Pecans and Walnuts).

VEGETABLES: Artichokes, Asparagus, Brassica (Cole) Vegetables (such as Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens and Rapini), Brassica Seed Production, Bulb Vegetables (such as Garlic, Leeks, Onions and Shallots), Cucurbit Vegetables (such as Cantaloupes, Cucumbers, Honeydew, Pumpkins, Squash and Watermelons), Fruiting Vegetables (such as Eggplant, Peppers and Tomatoes), Ginseng, Herbs and Spices (such as Anise (Fennel), Basil, Chive, Clove, Dill, Mint, Nutmeg, Parsley, Rosemary and Sage), Leafy Vegetables (such as Arugula, Celery, Cress, Endive, Lettuce, Radicchio, Rhubarb, Spinach, Swiss Chard), Legume Vegetables (such as all varieties of Beans and Peas, Garbanzos and Lentils), Okra, Root and Tuber Vegetables (such as Beets, Carrots, Ginger, Potatoes, Radishes, Sweet Potatoes, Turnips and Yams) and Root and Tuber Vegetable Seed Production.

NON-FOOD CROPS: Jojoba, Ornamentals and Turf (Sod and Seed production).

MIXING INSTRUCTIONS

Earthtrend Soil is water soluble and suitable for use in conventional liquid application systems.

Product Dilution: Thoroughly shake **Earthtrend Soil** and dilute in sufficient water to assure adequate and even coverage.

Dilution of water pH: Adjust alkaline dilution water (pH greater than 7) to pH 6 prior to adding **Earthtrend Soil**. Agitate the tank mixture during application and use within 12 hours after dilution. For best results use **Earthtrend Soil** with a full fertilization program.

COMPATIBILITY

Apply **Earthtrend Soil** at the end of the water application in sufficient water to penetrate into the root zone without excessive leaching into deeper soil. **Earthtrend Soil** may be tank mixed with herbicides, insecticides, fungicides, nematocides and fertilizers, or impregnated on granular fertilizers. Add **Earthtrend Soil** last to the spray tank mixes containing other fully diluted chemicals. Test the compatibility of the intended tank mixture before use.

For use with fertilizers containing a high phosphorous analysis, add one gallon of water to the spray tank for each three gallons of fertilizer, and add a buffer/compatibility agent prior to adding **Earthtrend Soil**. The following procedures may be helpful in the event of incompatibility:

- Predilute **Earthtrend Soil** in 5 gallons of water before adding to the spray tank.
- Increase the amount of water per acre to be applied.
- Add a buffer/compatibility agent to the spray tank.

Earthtrend Foliar Plant Growth Regulator APPLICATION TABLE

CROP	Early Seedling Vigor	Improved Root Growth	Stress Resistance	Uniformity/Improved Quality	Improved Fruit Size	Increased Fruit Number	Application Method	Use with Earthtrend Foliar	APPLICATION TIMING
Alfalfa	✓	✓	✓				B, C	O	At planting, apply Earthtrend Soil with starter fertilizer in 1st irrigation. Make additional applications with fertilizer in irrigation water 14 days after each cutting, or when new growth is initiated.
Artichokes, Asparagus	✓	✓	✓	✓	✓		B, C	O	Apply at planting and/or at the beginning of each new growth cycle. An additional application may be made 3 to 6 weeks after the first application.
Avocados, Bananas, Coffee, Figs, Guavas, Mangos, Papayas, Persimmons, Pineapple, Plantains	✓	✓	✓	✓	✓	✓	A, C	Y	Chemigate into the soil at initiation of root flush. Repeat applications at 30-day intervals during periods of active root growth. As a transplant treatment for seedlings or root stock, soak bare roots in Earthtrend Soil solution up to 6 hours before transplanting. Make a 2 nd application 30 days after first.

continued

Berry Fruits such as: Blueberries, Blackberries, Currants, Huckleberries, Loganberries, Raspberries	✓	✓	✓	✓	✓	✓	A, C	Y	Chemigate into the soil at initiation of root flush. Repeat application at 30-day intervals during periods of active root growth. As a transplant treatment for seedlings or root stock, soak bare roots in Earthtrend Soil solution up to 30 minutes before transplanting. Make a 2 nd application 30 days after first.
Brassica (Cole) Vegetables such as: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens, Rapini	✓	✓	✓	✓			A, B, C	O	Band 2 inches below seed, or rotovate into top 2 inches of soil with starter fertilizer at planting, or apply in 1st irrigation. Make a 2nd application 20 to 30 days after the 1st. For transplants, apply as a seedling soak or side-dress or chemigate with fertilizer within 7 days of transplanting. Make a 2 nd application 20 to 30 days after the first.
Bulb Vegetables such as: Garlic, Leeks, Onions (dry, bulb, green), Shallots	✓	✓	✓	✓	✓		B, C	O	Band 2 inches below the seed or sets with starter fertilizer or apply in first irrigation. Make a 2 nd application 30 days after the first.
Canola (Rape) and Cereal Grains such as: Barley, Corn, Oats, Rye, Sorghum (Milo), Wheat	✓	✓	✓	✓		✓	B, C, E	O	Apply as a seed treatment at planting. Or, band with starter fertilizer at planting. Or, sidedress or chemigate in between emergence and 4 th leaf stage.
Citrus Fruits such as: Grapefruit, Lemons, Limes, Mandarins, Oranges	✓	✓	✓	✓		✓	A, C	Y	Chemigate into soil at initiation of root flush. Repeat application at 30-day intervals during periods of active root growth. As transplant treatment for seedling or root stock, soak bare roots in Earthtrend Soil solution up to 30 minutes before transplanting.
Cotton	✓	✓	✓	✓			B, C	Y	Make 1 st application at planting. Make a 2 nd application with first side-dress.

continued

Cucurbit Vegetables such as: Cantaloupe, Cucumbers, Gourds, Honeydew, Muskmelons, Pumpkins, Squash (summer & winter), Watermelons	✓	✓	✓	✓	✓	✓	B, C	O	Apply preplant or in first fertilization after emergence. Make a 2nd application before layby or when plants begin to run.
Fruiting Vegetables such as: Eggplant, Ground Cherries, Peppers (all varieties), Tomatillo, Tomatoes	✓	✓	✓	✓	✓	✓	B, C	O	Apply at transplant, preplant, 1 st fertilization or 1 week after transplanting. Make a 2 nd and 3 rd application at 14 to 21 day intervals. For Roma Tomatoes , follow with an application of Earthtrend Foliar (See Earthtrend Foliar label for use directions).
Ginseng	✓	✓	✓	✓			B, C	O	Band 2 inches below seed, or rotovate into top 2 inches of soil with starter fertilizer or apply in 1 st irrigation. Make a 2 nd application 30 days after the 1 st . In the 2 nd year, apply at the initiation of growth in the spring.
Grapes (all varieties – dried, processed, wine, table)	✓	✓	✓	✓	✓	✓	A, C	O	Apply at initiation of each root flush. As a transplant treatment for new vines or root stock, soak bare roots in Earthtrend Soil solution up to 2 hours before transplanting.
Herbs and Spices such as: Anise, Basil, Chive, Clove, Dill, Mint, Nutmeg, Parsley, Rosemary, Sage	✓	✓	✓	✓	✓	✓	A, B, C	O	Apply transplant, preplant, at planting with a starter fertilizer or with 1 st fertilization after transplanting. Make a 2nd application 20-30 days after the 1 st application.
Hops	✓	✓	✓	✓		✓	B, C, E	Y	Apply as a seed treatment at planting. Or, band with starter fertilizer at planting. Or, chemigate in at emergence.
Jojoba	✓	✓	✓	✓	✓		B, C	O	Apply at preplant or planting; for established jojoba, apply twice a year at beginning of each growth cycle and again 10 to 15 days later.

continued

Kiwifruit	✓	✓	✓	✓	✓	✓	A, C	O	Chemigate into the soil at initiation of root flush. Repeat applications at 30-day intervals during periods of active root growth. As transplant treatment for seedlings, soak roots in Earthtrend Soil solution for 15-30 minutes before transplanting.
Leafy Vegetables such as: Arugula, Celery, Cress, Endive, Lettuce, Radicchio, Rhubarb, Spinach, Swiss Chard	✓	✓					B, C	O	Band 2 inches below seed, or rotate into top 2 inches of soil with starter fertilizer, or apply in 1 st irrigation. Make 2 nd application 20 days after the 1 st application. For transplant, apply as a seedling soak. Make a 2nd application 1 week after transplanting. Celery (Seeded): Band 2 inches below the seed. Reapply when seed row is visible and at 4 inches of growth. Celery (Transplants): Chemigate within 7 days after transplanting. Make 2 additional applications at 10-day intervals beginning at 3 inches of growth.
Legume Vegetables such as: Beans (all varieties), Garbanzos, Lentils, Peas (all varieties), Soybeans	✓	✓	✓	✓	✓	✓	B, C, E	Y	Apply preplant or at planting with fertilizer or as a seed treatment, or in first irrigation after planting. Or, apply as a side-dress when row is visible.
Okra	✓	✓	✓	✓	✓	✓	B, C	O	Apply as side-dress or chemigate in with first fertilization.
Ornamental Trees, Shrubs, Flowers	✓	✓	✓	✓			D	Y	Chemigate into the soil drench according to the schedule given in "Application Methods and Use Rates" section.
Peanuts	✓	✓	✓	✓			B	O	Apply at planting with a starter fertilizer or with first side-dress.
Pome Fruits such as: Apples, Crabapples, Pears, Quinces	✓	✓	✓	✓			A, C	Y	Chemigate into the soil at initiation of root flush. Repeat applications at 30-day intervals during periods of active root growth. As a transplant treatment for seedlings or root stock, soak bare roots in Earthtrend Soil solution up to 30 minutes before transplanting. To reduce alternate bearing, make a postharvest application.
Rice	✓	✓	✓	✓			B, C	Y	Make 1 st application prior to seeding or just prior to flooding the field. Make a 2 nd application any time before panicle initiation.

continued

Root and Tuber Vegetables such as: Beets, Carrots, Ginger, Horseradish, Potatoes, Potato Seed Production, Radishes, Sweet Potatoes, Turnips, Yams	✓	✓	✓	✓	✓	✓	A, B, C	O	Band 2 inches below the seed, seed piece or slip, or rotovate into the top 2 inches of soil with starter fertilizer, or apply with fertilizer in the first irrigation. Make a 2 nd application 20 days after the 1 st for short season crops or 20 to 30 days after the 1 st application for longer season crops. Potatoes: Side-dress or chemigate with fertilizer at tuber initiation or 40 to 6 inch rosette. Sweet Potatoes and Yams: Apply with starter fertilizer in transplant water.
Stone Fruits such as: Apricots, Cherries, Peaches, Plums, Nectarines	✓	✓	✓	✓	✓	✓	A, C	Y	Chemigate into the soil at initiation of root flush. Repeat applications at 30-day intervals during periods of active root growth. As a transplant treatment for seedlings or root stock, soak bare roots in Earthtrend Soil solution up to 30 minutes before transplanting.
Strawberries	✓	✓		✓	✓	✓	C	O	Chemigate through drip system at 21- to 30-day intervals beginning 7 days after transplanting. As a transplant treatment for seedlings, soak roots in Earthtrend Soil solution up to 30 minutes before transplanting.
Sugarcane	✓	✓	✓				A, C	O	Soak seed pieces for 20 seconds in solution before planting. Side-dress or chemigate in a 2 nd application with fertilizer at the 4-leaf stage.
Tree Nuts such as: Almonds, Cashews, Filberts (Hazelnuts), Pecans, Walnuts	✓	✓	✓	✓	✓		A, C	Y	Chemigate into the soil at initiation of root flush. Repeat applications at 30-day intervals during periods of active root growth. As a transplant treatment for seedlings or root stock, soak bare root in Earthtrend Soil solution up to 30 minutes before transplanting. To reduce alternate bearing, make a postharvest application.
Turf: Bermuda seed	✓	✓	✓	✓			B, C	O	Apply with fertilizer to initiate new growth.

*Some applications of **Earthtrend Soil** are improved when made in combination with **Earthtrend Foliar**.
Y = Yes; N = No; O = Optional.

APPLICATION METHODS

Apply Earthtrend Soil using the following methods:

- TRANSPLANT USE:** Apply **Earthtrend Soil** in the transplant water at rates stated in the APPLICATION TABLE or "Row Spacing Table", or use as a seedling soak or bare root soak prior to transplanting at 8 fluid ounces **Earthtrend Soil** per 100 gallons of water.
- PREPLANT OR AT PLANTING APPLICATION:** Broadcast spray **Earthtrend Soil** over the soil or seed furrow at 15-45 fluid ounces per acre and incorporate prior to planting. **Earthtrend Soil** may be banded directly into the seed furrow or with a starter fertilizer below or to the side of the seed row. For Cotton, Corn

and Sorghum (Milo) a minimum of 10 fluid ounces per acre may be applied. Alternatively, **Earthtrend Soil** may be chemigated into the seed row in the first irrigation, impregnated on dry fertilizers prior to preplant or side-dress applications. See ROW SPACING/RATE TABLE for rates.

- C. **POST-EMERGENT APPLICATIONS:** Knife **Earthtrend Soil** into the root zone after transplanting or thinning or when a seedling line is visible. Use ROW SPACING/RATE TABLE to calculate rates. **Earthtrend Soil** may be applied through sprinkler and drip irrigation systems. Time applications to allow penetration of **Earthtrend Soil** into the root zone without excessive leaching into deeper soil. See “Chemigation” section of this label for additional information.
- D. **NURSERY AND CONTAINER USE:** Apply **Earthtrend Soil** as a soil drench at the following dilutions and rates based on container size.

Container (gallons)	Dilution (pt/100 gallons)	Timing
1	1	Weekly
5	1	Bi-weekly
10 or more	1	3 weeks to monthly

- E. **SEED TREATMENTS:** Do not use treated seed for food, feed or oil purposes. Use 4 to 8 fluid ounces per acre of seed to be planted. Dilute **Earthtrend Soil** into seed treatment solution, and apply mixture directly onto the seed. Do not soak seeds in undiluted **Earthtrend Soil**. To determine optimal amount of **Earthtrend Soil** to use per pound of seed, please contact Westbridge Ag. Products at (800) 876-2767 Monday-Friday 9AM-5PM PT.
- HOPPER-BOX/OTHER AT PLANTING SEED TREATMENTS:** **Earthtrend Soil** may be applied directly onto seeds as a seed treatment immediately prior to or at planting. Treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time.
 - COMMERCIAL SEED TREATMENT:** An approved dye must be added before seeds may be sold in interstate commerce to distinguish treated seed, and to prevent inadvertent use for food, feed or oil purposes. Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal and State seed laws.

ROW SPACING/RATE TABLE

For many crops, **Earthtrend Soil** is applied to the seed furrow, as a side-dress, or through drip irrigation systems. Use 0.75 to 1.0 fluid ounce per 1,000 feet of row, or use the following table to determine application rate:

Row Spacing (Inches)	Earthtrend Soil (Fl. Oz./Acre)
40 or more	10 to 20
35	20 to 25
30	25 to 30
25	30 to 35
20 or less	35 to 45

HEAVY METAL INTERNET STATEMENT

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.appfco.org/metals.htm>.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store **Earthtrend Soil** in a cool place out of direct sunlight.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non-refillable container; do not reuse or refill this container. Triple rinse (or equivalent) container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling, if available; or reconditioning, if appropriate; or puncture and dispose of in a sanitary landfill; or, if allowed by State and local authorities, by incineration. If incinerated, stay out of smoke.

NOTICE OF WARRANTY

Westbridge warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Westbridge. To the fullest extent permitted by State law, Westbridge shall not be liable for consequential, special or indirect damages resulting from the use or handling of this product. Westbridge makes no warranties of merchantability or fitness for a particular purpose or any other express or implied warranty except as stated above.

Earthtrend Soil is manufactured by Westbridge Agricultural Products. TRIGGRR is a registered trademark of Westbridge Agricultural Products.

NOTES

Earthtrend Soil

PLANT GROWTH REGULATOR

Active Ingredient*:	
Cytokinin (as kinetin)	00.004%
Other Ingredients:	99.996%
Total	100.000%

* Contains Non-Plant Food Ingredients

** Based on Biological Activity

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for FIRST AID, PRECAUTIONARY STATEMENTS,
and DIRECTIONS FOR USE

SHAKE WELL BEFORE USING

CHEMIGATION: Refer to supplemental labeling entitled APPLICATION GUIDE for use directions for chemigation.

Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.

Refer to supplemental labeling entitled AGRICULTURAL USE REQUIREMENTS in the DIRECTIONS FOR USE section of the labeling for information about this standard.

EPA Reg. No. 51517-3 • EPA Est. No. 51517-CA-1

Net Contents: 1 U.S. Gallon

FIRST AID

IF INHALED:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.• Call a poison control center or doctor for further treatment advice.
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For more information call 1-800-876-2767 Monday-Friday 8-5 PT. After hours, weekends or holidays call 1-800-222-1222, the National Poison Control Center.

NOTICE OF WARRANTY - Westbridge warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of Westbridge. To the fullest extent permitted by State law, Westbridge shall not be liable for consequential, special or indirect damages resulting from the use or handling of this product. Westbridge makes no warranties of merchantability or fitness for a particular purpose or any other express or implied warranty except as stated above.

Manufactured by
 **Westbridge**
Agricultural Products

1260 Avenida Chelsea, Vista, CA 92081 USA